

Light quark masses and CKM matrix elements from lattice QCD

V. Lubicz^a

^aDip. di Fisica, Università di Roma Tre and INFN, Sezione di Roma Tre,
Via della Vasca Navale 84, I-00146 Roma, Italy

I give a brief overview of recent results from lattice QCD calculations which are relevant for the phenomenology of the Standard Model. I discuss, in particular, the lattice determination of light quark masses and the calculation of those hadronic quantities, such as semileptonic form factors, decay constants and B -parameters, which are of particular interest for the analysis of the CKM mixing matrix and the origin of CP violation.

1. LIGHT QUARK MASSES

Calculations of light quark masses are becoming, at present, one of the most important subject of investigation for lattice QCD. Indeed, in spite of their relevant phenomenological interest, light quark masses are still among the less known fundamental parameters of the Standard Model.

In the study of quark masses, the lattice method provides a unique approach: quark masses, defined as effective couplings renormalized at short distances, can be in fact determined on the lattice from non-perturbative calculations of hadronic quantities. An inspection of recent lattice results for quark masses suggests, however, that in such calculations a better understanding and quantification of the systematic errors is still an important requirement. To be specific, let us consider the lattice determinations of the strange quark mass, $\bar{m}_s$. A compilation of results for this mass, in the $\overline{\text{MS}}$ scheme, at the renormalization scale $\mu = 2 \text{ GeV}$, is shown in Table 1. All the results have been obtained in the quenched approximation. As can be seen from the table, although the lattice predictions all lie in the range $100 \lesssim \bar{m}_s \lesssim 160 \text{ MeV}$, however, within the errors quoted by the authors, the results obtained from different numerical studies are often in disagreement.

The main two sources of systematic errors, which are responsible for the discrepancies discussed above, are easily identified. The quark mass that is directly computed in lattice simulations is the (short-distance) bare lattice quark

Table 1

Lattice results for the strange quark mass $\bar{m}_s(2 \text{ GeV})$ in the $\overline{\text{MS}}$ scheme. We denote by $\mathcal{O}(a^2)$ and NPR the results obtained by using the non-perturbatively $\mathcal{O}(a)$ -improved action and non-perturbative renormalization respectively.

	Yr	$\bar{m}_s(2 \text{ GeV})/\text{MeV}$	
APE [1]	94	128 ± 18	
LANL [2]	96	$100 \pm 21 \pm 10$	
FNAL [3]	96	95 ± 16	
APE [4]	96	122 ± 20	
SESAM [5]	97	166 ± 15	
CP-PACS [6]	97	135 ± 7	(from m_ϕ)
	97	111 ± 4	(from m_K)
JLQCD [7]	97	97 ± 9	
QCDSF [8]	97	112 ± 5	$\mathcal{O}(a^2)$
APETOV [9]	97	111 ± 15	$\mathcal{O}(a^2)$
APE [10]	98	$130 \pm 2 \pm 18$	NPR
APE [11]	98	121 ± 13	$\mathcal{O}(a^2) + \text{NPR}$

mass $m(a)$, where a is the lattice spacing. This quantity is typically determined by fixing, to its experimental value, the mass of a hadron containing a quark with the same flavour. Since the non-perturbative lattice calculation of hadronic quantities is affected by discretization errors (in general of $\mathcal{O}(a)$), the same systematic errors will also propagate into the final determination of the quark masses.

The second step in these calculations consists in relating the bare quark mass $m(a)$ to the renormalized mass $m(\mu)$, in a given renormalization scheme. The connection is provided by a multiplicative renormalization constant, $m(\mu) =$

$Z_m(a\mu) m(a)$. The perturbative, renormalization group improved expression of $Z_m(a\mu)$, at the NLO, has been given in ref. [1]. Clearly, this matching introduces higher orders perturbative errors, which in turn affect the determination of the quark mass.

It is a remarkable fact that, at present, both discretization and higher orders perturbative errors, in the lattice determination of light quark masses, can be (and in fact have been) drastically reduced. The way to deal with discretization errors has been suggested by Symanzik [12] and consists in improving the lattice action and operators. This program has been realized perturbatively in [13,14] and, more recently, carried out at the non-perturbative level in [15]-[17]. The discretization errors are reduced to $\mathcal{O}(a^2)$ in the chiral limit. The residual cutoff effects are expected to affect the determination of light quark masses by less than 1% [11].

Higher orders perturbative errors in the evaluation of the quark mass renormalization constant are eliminated by adopting the non-perturbative renormalization prescription proposed in ref. [18]. The idea consists on imposing the renormalization conditions directly on non-perturbatively calculated correlation functions between external off-shell quark states. This prescription defines the mass in a specific scheme, called RI-MOM scheme. Lattice calculations in this scheme are thus completely non-perturbative. Perturbation theory only enters if one wants to convert the result to the mass in the $\overline{\text{MS}}$ scheme. In this step (which is not necessary anyway) higher orders perturbative errors are expected to be negligible, since the relevant conversion factor is presently known at the N²LO [19].

Among the lattice predictions for the strange quark mass (see Table 1), the smallest values, corresponding to $\overline{m}_s \lesssim 100$ MeV, have been obtained in refs. [2,3,7] as a result of an extrapolation to the continuum limit, $a \rightarrow 0$. However, the $\mathcal{O}(a)$ -improved calculations of refs. [8,9,11], in which discretization effects are expected to be negligible, predict larger values, thus casting some doubts on the reliability of previous extrapolations.

In ref. [11] the first lattice calculation of light

quark masses, which combined both the use of a non-perturbatively improved action and the non-perturbative renormalization technique, has been presented. As already noted in [10], the non-perturbative calculation of renormalization constants removes the discrepancies, observed in previous calculations, between the results obtained by using different definitions of the bare quark mass (based on the vector and the axial-vector Ward identities respectively). It also gives values of the quark masses larger than those obtained by computing the renormalization constants in perturbation theory. The relevant renormalization constants which enter in the lattice calculations of the quark masses are those of the scalar and pseudoscalar densities, Z_S and Z_P . By comparing the values of these constants obtained by using the non-perturbative method with the predictions of one-loop (boosted) perturbation theory it is found that the differences can be larger than 10% and 30% in the case of Z_S and Z_P respectively. Clearly, this error is introduced in the determination of the quark masses, when the renormalization constants are evaluated in perturbation theory. As an example of a non-perturbative determination of renormalization constants, we show in Figure 1 the values of Z_S as a function of the renormalization scale μ [11]. It is reassuring to find that this behaviour results in excellent agreement with the dependence predicted by the renormalization group equation.

As a final estimate of the average light ($\overline{m}_l = (\overline{m}_u + \overline{m}_d)/2$) and strange quark masses, in the $\overline{\text{MS}}$ scheme at the scale $\mu = 2$ GeV, I quote the results of ref. [11]:

$$\begin{aligned} \overline{m}_l^{NLO} &= 4.9(4) \text{ MeV} \\ \overline{m}_s^{NLO} &= 121(13) \text{ MeV} \end{aligned} \quad (1)$$

and

$$\begin{aligned} \overline{m}_l^{N^2LO} &= 4.5(4) \text{ MeV} \\ \overline{m}_s^{N^2LO} &= 111(12) \text{ MeV} \end{aligned} \quad (2)$$

at the NLO and N²LO respectively. In these estimates discretization and higher orders perturbative errors are expected to be negligible, and the remaining uncertainty is mainly due to the quenched approximation. The values in eq. (1)




Figure 1. Z_S , obtained by using the non-perturbative method, as a function of the renormalization scale. The dashed and solid curves represent the solutions of the renormalization group equations at the LO and NLO respectively.

are also in good agreement with recent results from QCD sum rules [20].

2. CKM MATRIX ELEMENTS

In the analysis of the CKM mixing matrix, a particularly interesting issue is represented by the study of the unitarity triangle in the ρ - η plane. Indeed, a non trivial shape of this triangle is the signature of a CKM source of CP violation.

At present, the phenomenological analysis of the unitarity triangle is constrained (mainly) by the existing measurements of $B_{d,s}^0 - \bar{B}_{d,s}^0$ mixings, $K^0 - \bar{K}^0$ mixing and semileptonic $b \rightarrow u$ transitions. Once these measurements are compared with the theoretical predictions, constraints on the ρ - η parameters are derived. The status of the art is shown in Figure 2, from ref. [21]. The theoretical predictions are based on the non-perturbative evaluation of some relevant hadronic matrix elements, which, in turn, are parameterized in terms of form factors, decay constants and B -parameters. The lattice method provide a reliable approach to perform such calculations. In the following, I will present a short compilation of these lattice results.

2.1. V_{ub} from $B \rightarrow \pi, \rho$ semileptonic decays

The determination of the V_{ub} mixing angle is allowed by the analysis of $B \rightarrow \pi, \rho$ semileptonic




Figure 2. The unitarity triangle in the $\bar{\rho}$ - $\bar{\eta}$ plane and the constraints derived from Δm_d , $\Delta m_s/\Delta m_d$, $|\epsilon_K|$ and $|V_{ub}|/|V_{cb}|$.

decays. The reliability of lattice calculations in the study of heavy mesons semileptonic decays is provided by the analysis of D -meson decays, $D \rightarrow K, K^* l \nu$. In this processes the relevant mixing angle V_{cs} is well constrained by the unitarity of the CKM matrix, so that the theoretical predictions can be compared with the experimental measurements. The comparison is shown in Figure 3, where the lattice results for the four relevant form factors, at zero momentum transfer, are presented together with their experimental values [29]. A summary of these results is also given in Table 2.




Figure 3. Lattice results for the $D \rightarrow K, K^* l \nu$ form factors, at zero momentum transfer, [22]-[28]. The horizontal band indicates the present experimental average.

For the $B \rightarrow \pi, \rho$ semileptonic decays, four different lattice groups have presented results so far

Table 2

Summary of lattice and experimental results for $D \rightarrow K, K^*$ semileptonic decays form factors at zero momentum transfer. The results for $B \rightarrow \pi, \rho$ semileptonic decays, from ref. [30], are also presented.

	$D \rightarrow K, K^*$		$B \rightarrow \pi, \rho$
	Lattice	Exp.	Lattice
$f_+(0)$	0.72(7)	0.76(3)	0.27(11)
$V(0)$	1.14(18)	1.07(9)	$0.35^{(+6)}_{(-5)}$
$A_1(0)$	0.64(8)	0.58(3)	$0.27^{(+5)}_{(-4)}$
$A_2(0)$	0.53(13)	0.41(5)	$0.26^{(+5)}_{(-3)}$

[24,26,28,30]. These are in quite good agreement one to each other. The more accurate determination of form factors has been obtained in ref. [30], from a combined analysis of B -meson semileptonic decays and radiative $B \rightarrow K^* \gamma$ decays. The results are shown in Table 2.

2.2. $\hat{B}_K$, the kaon bag parameter

The bag parameter $\hat{B}_K$ is the relevant hadronic quantity which enters in the calculation of ϵ_K . A recent compilation of lattice results gives, for the NLO renormalization group invariant parameter, the estimate [31]:

$$\hat{B}_K = 0.86(6)(6) \quad (3)$$

Notice that, in present analysis of the unitarity triangle, four constraints are used to determine the values of two parameters, ρ and η (see Figure 2). Thus, one of the constraints can also be removed, and the corresponding quantity can be determined together with ρ and η . In this way, in ref. [21] they obtain the value

$$\hat{B}_K = 0.82^{+0.45}_{-0.24} \quad (4)$$

in very good agreement with the lattice determination of eq. (3). This result also implies that significantly lower estimates of $\hat{B}_K$, such as those obtained by using the QCD hadronic duality approach ($\hat{B}_K = 0.39 \pm 0.10$) [32] or using the $SU(3)$ symmetry and PCAC ($\hat{B}_K = 1/3$) [33] are presently strongly disfavoured by the experiments.

Table 3

Summary of lattice results for D and B -meson decay constant and B parameters

$f_D/(\text{MeV})$	$f_{D_s}/(\text{MeV})$	$f_B/(\text{MeV})$
200(30)	220(30)	170(35)
$f_B \sqrt{B_B}/(\text{MeV})$	f_{B_s}/f_B	B_{B_s}/B_B
201(42)	1.14(8)	1.00(3)

2.3. Δm_d and Δm_s from B -meson decay constants and B -parameters

The theoretical estimates of the B -meson mass differences, Δm_d and Δm_s , can be expressed in terms of the pseudoscalar decay constants, $f_{B_{d,s}}$, and B -parameters. A recent compilation of lattice results for these quantities has been presented in ref. [34] and it is shown in Table 3. Notice that the lattice value for f_{D_s} , which has been predicted well before the first experimental measurement, it is in good agreement with the present experimental average, $f_{D_s} = 243 \pm 36 \text{ MeV}$ [21]. The lattice results for this quantity have always been stable in the time. This is shown in Figure 4, where these results, obtained over a period of 10 years, are presented together with the current experimental value. The last point in




Figure 4. Lattice results for the pseudoscalar decay constant f_{D_s} . The horizontal band indicates the present experimental average. We refer to ref. [34] for a compilation of results and corresponding references.

the figure represents the preliminary result of the first $\mathcal{O}(a)$ -improved lattice calculation of the heavy mesons decay constants. It corresponds to $f_{D_s} = (230 \pm 12) \text{ MeV}$ [35].

The lattice determinations of the B -meson decay constant, f_B , still represent a genuine prediction, since this constant has not been mea-

sured yet in the experiments. From the (overconstrained) fit of the Standard Model, in ref. [21] the value

$$f_B \sqrt{B_B} = 213(21)(20) \text{ MeV} \quad (5)$$

is obtained, in good agreement with the lattice determination (see Table 3). An independent determination of f_B can be also derived by combining the lattice determination of the ratio f_B/f_{D_s} , in which many of the systematic errors are expected to cancel, with the experimental measurement of f_{D_s} . Two recent lattice calculations give:

$$f_B/f_{D_s} = \begin{cases} 0.77(8) & \text{APE [35]} \\ 0.75(5)(7) & \text{MILC [36]} \end{cases} \quad (6)$$

From these values I find:

$$f_B = (185 \pm 27 \pm 19) \text{ MeV} \quad (7)$$

where the first error comes from the experimental uncertainty on f_{D_s} and the second one from the theoretical uncertainty on the ratio f_B/f_{D_s} . Eq. (7) is also in good agreement with the direct determination of f_B given in Table 3.

REFERENCES

1. C.R. Allton *et al.*, Nucl.Phys. **B431** (1994) 667.
2. R. Gupta and T. Bhattacharya, Phys.Rev. **D55** (1997) 7203.
3. B.J. Gough *et al.*, Phys.Rev.Lett. **79** (1997) 1622.
4. C.R. Allton, V. Gimenez, L. Giusti and F. Rapuano, Nucl.Phys. **B489** (1997) 427.
5. N. Eicker *et al.*, SESAM Collaboration, Phys.Lett. **B407** (1997) 290.
6. S. Aoki *et al.*, CP-PACS Collaboration, Nucl.Phys. (Proc.Suppl.) **63** (1998) 161; Nucl.Phys. (Proc.Suppl.) **60A** (1998) 14.
7. S. Aoki *et al.*, JLQCD Collaboration, Nucl.Phys. (Proc.Suppl.) **63** (1998) 275.
8. M. Gockeler *et al.*, Phys.Rev. **D57** (1998) 5562.
9. A. Cucchieri *et al.*, Phys.Lett. **B422** (1998) 212.
10. V. Gimenez *et al.*, EDINBURGH-97-15. hep-lat/9801028
11. D. Becirevic *et al.*, LPTHE-ORSAY-98-37. hep-lat/9807046.
12. K. Symanzik, Nucl.Phys. **B226** (1983) 187 and 205.
13. B. Sheikholeslami and R. Wohlert, Nucl.Phys. **B259** (1985) 572.
14. G. Heatlie *et al.*, Nucl.Phys. **B352** (1991) 266.
15. M.Lüscher *et al.*, Nucl.Phys. **B478** (1996) 365.
16. M.Lüscher *et al.*, Nucl.Phys. **B491** (1997) 323.
17. M.Lüscher *et al.*, Nucl.Phys. **B491** (1997) 344.
18. G. Martinelli *et al.*, Nucl.Phys. **B445** (1995) 81.
19. E. Franco and V. Lubicz, ROME-I-1198-98, to appear in Nucl.Phys.B. hep-ph/9803491.
20. C.A.Dominguez, these proceedings.
21. P. Paganini *et al.*, DELPHI-97-137, hep-ph/9711261; F. Parodi *et al.*, hep-ph/9802289.
22. V. Lubicz *et al.*, Nucl.Phys. **B356** (1991) 301; V. Lubicz *et al.*, Phys.Lett. **B274** (1992) 415.
23. C.W. Bernard *et al.*, Phys.Rev. **D43** (1991) 2140; Phys.Rev. **D45** (1992) 869.
24. A. Abada *et al.*, ELC Collaboration, Nucl.Phys. **B416** (1994) 675.
25. K.C. Bowler *et al.*, UKQCD Collaboration, Phys.Rev. **D51** (1995) 4905.
26. C.R. Allton *et al.*, APE Collaboration, Phys. Lett. **B345** (1995) 513.
27. T. Bhattacharya and R. Gupta, Nucl.Phys. B(Proc.Suppl.) **47** (1996) 481.
28. S. Gusken, private communication as quoted in ref.[34].
29. A. Ryd, presented at the 1997 Heavy Flavour Symposium, Santa Barbara, California.
30. L. Del Debbio *et al.*, Phys.Lett. **B416** (1998) 392.
31. R. Gupta, LAUR-98-271. hep-ph/9801412.
32. A. Pich and E. de Rafael, Phys.Lett. **B158** (1985) 477; J. Prades *et al.*, Z.Phys. **C51** (1991) 287.
33. J.F. Donoghue, E. Golowich and B.R. Holstein, Phys.Lett. **B119** (1982) 412.
34. J.M. Flynn and C.T. Sachrajda, to appear in Heavy Flavours (2nd ed.), ed. by A.J. Buras and M. Linder (World Scientific, Singapore). hep-lat/9710057.
35. D. Becirevic *et al.*, in preparation.
36. C. Bernard *et al.*, MILC Collaboration, FSU-SCRI-98-65. hep-ph/9806412.